

**EMBARGOED UNTIL FRIDAY 26 FEBRUARY, 2021, 10am GMT**

Media release, 26 February 2021

## **SARS-CoV-2 mutations in competition**

**How dangerous are new mutations of the SARS-CoV-2 virus? An international team involving researchers from the Institute of Virology and Immunology (IVI) of the Federal Food Safety and Veterinary Office BLV and the University of Bern (Switzerland), the Centers for Disease Control and Prevention (USA), and the Friedrich-Loeffler Institute (Germany), has developed an approach that can accurately assess the transmissibility of new virus mutants.**

Prior to the emergence of new mutants of the coronavirus, such as the British variant B.1.1.7, the SARS-CoV-2 variant named D614G was the most widespread variant worldwide. However, D614G had already mutated from the original SARS-CoV-2 pathogen that triggered the pandemic. An international team including researchers from Bern has now been able to demonstrate in both the laboratory and in animal models why the D614G variant was able to gain the upper hand over the original SARS-CoV-2 virus. "Our approach also allows us to characterize emerging mutations such as the British variant B.1.1.7 better and quicker," says Volker Thiel of the Institute of Virology and Immunology (IVI), one of the four lead authors of the study. The findings are extremely important for assessing the risk of new mutants running rampant, as they show how a fitness advantage of virus variants can lead to higher transmission. First results were released earlier allowing for scientific discussion on what is known as a preprint server. The results of the study have now been published in full in *Nature*.

The D614G variant carries a mutation in the spike protein that makes it easier for the virus to dock onto human cells. The researchers at IVI and in David E. Wentworth's laboratory at the Centers for Disease Control and Prevention in Atlanta (USA) first demonstrated in human cell cultures from the upper respiratory tract, as well as from the nose, that the D614G variant binds more strongly and also replicates faster than the original virus. In the animal model with mice, these findings were also confirmed in vivo, i.e., in the living organism. These experiments were then carried out at the IVI in Charaf Benarafa's group.

### **The new mutation clearly prevails**

The spread of SARS-CoV-2 viruses can be studied better in other animals rather than mice. Hamsters and ferrets are well established in infection research and are especially suitable animal models. To measure transmissibility, a mixture of equal parts of the original version of the SARS-

CoV-2 virus and the D614G variant was applied into the nose of each animal under light anesthesia. After one day, individual animals were rehoused with another healthy animal of the same species, and the transmission of the two variants measured. Thus, the infectivity of the two viruses could be observed in direct competition with each other. The experiment was repeated with six pairs of animals in total. In virtually all newly infected animals, the proportion of transmitted SARS-CoV-2 viruses was almost completely dominated by the D614G variant early on. The differentiation of the variants was carried out using the latest sequencing technology and PCR techniques by Martin Beer's team at the Friedrich Loeffler Institute, Federal Research Institute for Animal Health, in Greifswald-Insel Riems (D). "Our study stands out because we were able to clearly discern the more efficient transmission of the mutated variant in direct comparison with the original variant," says Volker Thiel.

### **A fitness test for further mutations**

This approach can even be used to test any single mutation or a specific combination of mutations that are present in a number of currently circulating viral variants. The IVI relies on a [cloning technique developed in Bern a year ago](#), in which SARS-CoV-2 viruses can be exactly reproduced in the laboratory. The British virus, for example, is known to have not just one but often more than 14 mutations, eight of which occur in the spike protein. Thus, with the help of the cloning technique, any number of mutations of variants can be reproduced and used to compete against each other in the established cell cultures and animal models. The results show how single mutations affect the fitness and transmissibility of new variants. "Our testing strategy allows us to rapidly examine why other, newly emerging virus variants have become established," says Volker Thiel.

Similar research projects on infectious pathogens could also be carried out in the future at the newly established [Multidisciplinary Center for Infectious Diseases and Immunity \(MCIDI\)](#) at the University of Bern.

The study was financially supported by the Swiss National Science Foundation SNF, the European Commission, the German Federal Ministry of Education and Research, and the U.S. Department of Health & Human Services, National Institutes of Health (NIH) / Institute of Allergy and Infectious Diseases (NIAID).

### **Institute of Virology and Immunology IVI**

The Institute of Virology and Immunology (IVI) with locations in Mitholz and Bern is a research institution affiliated with the Federal Food Safety and Veterinary Office FSVO with participation of the University of Bern. It is the only high-security laboratory in Switzerland where highly contagious animal diseases (such as foot-and-mouth disease or swine fever) can be diagnosed and researched. In line with the "one health" concept, one research focus is on viral infections that pose a threat to animals and humans (zoonoses).

[Further information](#)

### **Publication details:**

Bin Zhou, Tran Thi Nhu Thao, Donata Hoffmann, Adriano Taddeo et al: *SARS-CoV-2 spike 614G variant confers enhanced replication and transmission*. Nature, February 25, 2021, [doi: xy./LINK](#)

**Contact:** Prof. Dr. Volker Thiel, Institute of Virology and Immunology (IVI), Federal Food Safety and Veterinary Office FSVO, and University of Bern.

Phone +41 31 631 2413 / [volker.thiel@vetsuisse.unibe.ch](mailto:volker.thiel@vetsuisse.unibe.ch)